

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Interactions of AtRGL1, a Negative Regulator of Gibberellic Acid Signalling

A thesis presented in partial fulfilment of the requirements for the degree of

**Master of Science
in
Biochemistry**

**Massey University, Palmerston North
New Zealand.**

**David John Sheerin
2005**

Abstract

Arabidopsis thaliana AtRGL1 (repressor of *ga1-3* like-1) is a negative regulator of the signal transduction pathway of the plant hormone gibberellin. AtRGL1 belongs to the DELLA subfamily within the GRAS family of plant regulatory proteins. There are four other DELLA proteins, including AtRGA (repressor of *ga1-3*) and AtRGL2, encoded by the *A. thaliana* genome. Previous studies provided evidence that the DELLA proteins are nuclear localised and are functionally divided into N- and C-terminal domains. The N-terminal domain perceives the gibberellin signal, while the C-terminal domain functions as a negative regulator of transcription and also as a possible dimerisation domain. Previous studies have also shown that *AtRGA*, *AtRGL1*, and *AtRGL2* function together in the regulation of the development of the inflorescence and that *AtRGL1* is primarily expressed in this tissue.

To investigate how DELLA proteins function in gibberellin signalling, I sought plant proteins that interact with AtRGL1. Two proteins, p24 (24 kDa) and p64 (64 kDa), were isolated from wild-type plant nuclear extracts by affinity to the N-terminal 121 amino acid residues of AtRGL1. The identity of these two proteins remains to be established. To investigate the interactions of the C-terminal domain of AtRGL1 an anti-AtRGL1 polyclonal antiserum was developed for co-immunoprecipitation experiments. However, AtRGL1 was not detectable in plant nuclear extracts from the inflorescence of wild-type plants, precluding this approach.

The possibility of DELLA protein dimerisation was also investigated using AtRGA, AtRGL1, and AtRGL2 in yeast 2-hybrid experiments. Yeast 2-hybrid protein interaction results suggest that AtRGA, AtRGL1, and AtRGL2 do not form homo- or hetero-dimers. Complexities encountered with this approach could make these results invalid, so these interactions require further investigation.

Acknowledgments

I would like to thank my supervisors Dr Jasna Rakonjac and Dr Toshi Foster for their guidance and support throughout this thesis. I would also like to thank all the people in the Helipad lab who have helped me through my research. I would also like to like to thank the people at the HortResearch Palmerston North plant immunology lab, who are lead by Dr William Jones, for supplying materials and advice. Thank you also to my friends and family for their support throughout the past two years.

This research was supported by a fellowship and funding from HortResearch.

Table of Contents

Abstract	ii
Acknowledgments	iii
Table of contents	iv
List of Figures	vii
List of Tables	viii
List of Abbreviations	ix
Note on nomenclature	xii
 Chapter 1. Introduction	
1.1. Introduction to plant hormones	1
1.2. Gibberellins act through a signal transduction pathway	2
1.2.1. Production of gibberellic acids	5
1.2.2. Cell perception of gibberellic acids	5
1.2.3. Cytoplasmic secondary messengers	7
1.2.4. Nuclear molecular switch	9
1.2.5. Gene regulation and cell responses	12
1.2.6. Regulation of gibberellin responses by other plant hormones	14
1.3. Organisation of DELLA proteins	16
1.4. Degradation of DELLA proteins	17
1.5. Developmental functions of DELLA proteins	22
1.6. Speculative gibberellic acid signalling pathway	25
1.7. How do DELLA proteins function?	27
1.7.1. Hypothesis	28
1.7.2. Aim 1	28
1.7.3. Aim 2	28
 Chapter 2. Materials and methods	
2.1. Materials and reagents	29
2.2. General methods	29
2.2.1. Molecular biology	29
2.2.2. Protein electrophoresis	29

2.2.3 Western blotting	30
2.3. Bacterial strains and culture conditions	30
2.4. Plant material and growth conditions	30
2.5. Yeast strains and culture conditions	31
2.6. Bacterial recombinant protein expression	31
2.6.1. Purification of MBP-LacZ	32
2.6.2. Purification of MBP-AtRGL1 ⁽¹⁻¹²¹⁾	32
2.6.3 Purification of AtRGL1 ⁽¹⁻¹²¹⁾ -TrxA	33
2.7. Extraction of plant nuclear proteins	34
2.8. Affinity purification	36
2.9. Mass spectroscopy	36
2.10. Anti-AtRGL1 polyclonal antibodies	36
2.11. Cloning of full length DELLA genes	37
2.12. Yeast 2-hybrid experiments	40
2.12.1. <i>HIS3</i> reporter gene assay	41
2.12.2. <i>LACZ</i> reporter gene assay	41
 Chapter 3. Results	
3.1. Affinity purification of AtRGL1-interacting proteins	47
3.1.1. Expression and purification of the AtRGL1 'bait' protein	49
3.1.2. Preparation of plant nuclear extract	51
3.1.3. Affinity purification of AtRGL1-interacting proteins	53
3.2. Antibody characterisation	60
3.2.1. Characterisation of an anti-DELLA peptide monoclonal antibody, 5E1	60
3.2.2. Production of an anti-AtRGL1 polyclonal antibody	63
3.2.3. Characterisation of the capacity of anti-AtRGL1 ⁽¹⁻¹²¹⁾ polyclonal antibodies to immunoprecipitate the antigen from solution	65
3.3. Analysis of DELLA-DELLA protein interactions	67
3.3.1. Expression of DELLA proteins in <i>Saccharomyces cerevisiae</i>	69
3.3.2. Yeast 2-hybrid analysis of DELLA-DELLA protein interactions	72

Chapter 4. Discussion	
4.1. Isolation of AtRGL1-interacting proteins	77
4.2. Investigation of DELLA-DELLA protein interactions	80
4.3. Future identification of AtRGL1-interacting proteins	83
4.4. Plans for immunoprecipitation of AtRGL1	84
4.5. Further investigation of yeast 2-hybrid experiments	87
Chapter 5. Conclusions	88
References	90

List of Figures

Figure 1-1 Phenotypes of <i>A. thaliana</i> GA signalling mutants.	3
Figure 1-2 Gibberellic acid signal pathway model.	4
Figure 1-3 DELLA protein domain organisation.	15
Figure 1-4 Structure and function of an SCF E3 ubiquitin ligase complex.	19
Figure 1-5 Detailed speculative gibberellic acid signal pathway model.	26
Figure 3-1 Schematic of co-affinity and affinity purification procedures.	48
Figure 3-2 MBP-AtRGL1 ⁽¹⁻¹²¹⁾ fusion protein purified from <i>E. coli</i> .	50
Figure 3-3 Soluble plant nuclear extract.	52
Figure 3-4 Affinity purification of MBP-AtRGL1 ⁽¹⁻¹²¹⁾ -interacting proteins from whole plant nuclear extract.	55
Figure 3-5 Affinity purification of MBP-AtRGL1 ⁽¹⁻¹²¹⁾ -interacting proteins from inflorescence nuclear extract.	57
Figure 3-6 Schematic of immunoprecipitation.	61
Figure 3-7 Western blot analysis of plant nuclear extract using anti-DELLA antibodies.	62
Figure 3-8 Fractionation and determination of the specificity of anti-MBP-AtRGL1 ⁽¹⁻¹²¹⁾ antiserum.	64
Figure 3-9 Immunoprecipitation of AtRGL1 ⁽¹⁻¹²¹⁾ -TrxA using purified anti-AtRGL1 ⁽¹⁻¹²¹⁾ polyclonal antibodies.	66
Figure 3-10 Schematic of yeast 2-hybrid analysis of DELLA-DELLA protein interactions.	68
Figure 3-11 Western blot detection of AD-DELLA fusions expressed in yeast.	70
Figure 3-12 Western blot detection of AD- and BD-DELLA protein fusions expressed in yeast.	71
Figure 3-13 <i>LACZ</i> reporter gene analysis of DELLA-DELLA protein yeast 2-hybrid interactions.	75
Figure 4-1 Schematic of possible DELLA-DELLA dimer interaction with reporter gene promoters.	82

List of Tables

Table 1. Plasmids	42
Table 2. Bacterial strains	43
Table 3. Primers	44
Table 4. Yeast strains	45
Table 5. Antibodies	46
Table 6. <i>HIS3</i> reporter gene assay of DELLA-DELLA protein interactions.	74

Abbreviations

3-AT	3-amino-1,2,4-triazole
ABA	Absciscic acid
AD	GAL4 activation domain
AMP	Ampicillin
AmSO ₄	Ammonium sulfate
AP	Alkaline phosphatase
AtASK1	<i>A. thaliana</i> Arabidopsis Skp like-1
AtGAI	<i>A. thaliana</i> Gibberellic acid insensitive
AtGAMYB33	<i>A. thaliana</i> Gibberellic acid induced MYB-33
AtGAMYB65	<i>A. thaliana</i> Gibberellic acid induced MYB-65
AtGCR1	<i>A. thaliana</i> G-protein coupled receptor-1
AtGPA	<i>A. thaliana</i> G-protein α subunit
AtHDA19	<i>A. thaliana</i> Histone deacetylase-19
AtLFY	<i>A. thaliana</i> Leafy
AtLRP	<i>A. thaliana</i> Lateral root primordium
AtPKL	<i>A. thaliana</i> Pickle
Atrga Δ 17	<i>A. thaliana</i> Repressor of <i>gal-3</i> , deletion of 17 amino acid DELLA motif
AtRGL1	<i>A. thaliana</i> Repressor of <i>gal-3-1</i>
Atrgl1 Δ 17	<i>A. thaliana</i> Repressor of <i>gal-3-1</i> , deletion of 17 amino acid DELLA motif
AtRGL2	<i>A. thaliana</i> Repressor of <i>gal-3-2</i>
AtRGL3	<i>A. thaliana</i> Repressor of <i>gal-3-3</i>
AtSHI	<i>A. thaliana</i> Short internodes
AtSLY1	<i>A. thaliana</i> Sleepy-1
AtSLY2	<i>A. thaliana</i> Sleepy-2
AtSNY	<i>A. thaliana</i> Sneezy (AtSLY2)
AtSPY	<i>A. thaliana</i> Spindly
BD	GAL4 DNA binding domain
BnSCL1	<i>B. napus</i> Scarecrow like-1
BSA	Bovine serum albumin
cDNA	complementary deoxyribonucleic acid
cGMP	Cyclic guanosine monophosphate
c-Myc	Mammalian c-Myc oncogene epitope

Co-IP	Co-immunoprecipitation
Col-0	Columbia-0
DAPI	4',6-Diamidino-2-phenylindole
DEAE	Diethylaminoethyl
DMSO	Dimethyl sulphoxide
DNA	Deoxyribonucleic acid
DTT	1,4-Dithiothreitol
EDTA	Ethylenediaminetetraacetic acid
GA	Gibberellic acid
GAL4	GAL4 transcription factor
GARE	Gibberellic acid responsive element
GFP	Green Fluorescent Protein
HA	Influenza hemagglutinin epitope
HDAC	Histone deacetylase
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HGP	Heterotrimeric G-protein
HRP	Horseradish peroxidase
HvCDPK1	<i>H. vulgare</i> Calcium dependent protein kinase-1
HvGAMYB	<i>H. vulgare</i> Gibberellic acid induced MYB
HvHSIMYB	<i>H. vulgare</i> Spindly interacting MYB
HvHSINAC	<i>H. vulgare</i> Spindly interacting NAC
HvSAD	<i>H. vulgare</i> Sad
HvSLN1	<i>H. vulgare</i> Slender-1
HvSPY	<i>H. vulgare</i> Spindly
IgG	Immunoglobulin G
Imidazole	1,3-Diaza-2,4-cyclopentadiene
IPTG	Isopropylthio- β -D-galactoside
KAN	Kanamycin
Ler-0	Landsberg <i>erecta</i> -0
MALDI TOF	Matrix assisted laser desorption ionisation time of flight
MBP	Maltose-binding protein
miR159	Micro ribonucleic acid 159
miRNA	Micro ribonucleic acid
mRNA	Messenger ribonucleic acid

MWCO	Molecular weight cut-off
NHS	<i>N</i> -hydroxysuccinimide
NLS	Nuclear localisation signal
<i>O</i> -GlcNAc	<i>O</i> -linked <i>N</i> -acetyl glucosamine
ONPG	<i>O</i> -Nitrophenol-galactoside
OsD1	<i>O. sativa</i> Dwarf-1
OsDOF3	<i>O. sativa</i> DNA binding with one finger-3
OsGID2	<i>O. sativa</i> Gibberellic acid insensitive dwarf-2
OsSLR1	<i>O. sativa</i> Slender rice-1
p24	24 kDa AtRGL1-interacting protein
p53	Murine p53
p64	64 kDa AtRGL1-interacting protein
PAGE	Poly-acrylamide gel electrophoresis
PBS	Phosphate buffered saline
PCR	Polymerase chain reaction
PEG	Poly-ethylene glycol
PMSF	Phenylmethanesulphonylfluoride
SCF	Skp-Cullin-F-box
SD	Synthetic dropout
SDS	Sodium dodecyl sulfate
StPHOR1	<i>S. tuberosum</i> Photoperiod-responsive-1
T antigen	SV40 large T antigen
TBS	Tris-buffered saline
TBST	Tris-buffered saline Tween-20
Tris	Tris (hydroxymethyl) aminomethane
TrxA	Thyroxine domain A
UV	Ultra violet

Note on Nomenclature

Gene names and symbols are written in *italics*, and proteins are in standard text. Wild-type genes and products are in uppercase, mutants are in lower case. For plant species, the two letters preceding a gene/protein refer to the species. At, *Arabidopsis thaliana*. St, *Solanum tuberosum*. Hv, *Hordeum vulgare*. Os, *Oryza sativa*. Bn, *Brassica napus*.